



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

For the *Accipiter velox rufilatus* Ridgway (1888) an older name is found in *Nisus pacificus* Lesson (1845), based on specimens from the western coast of Mexico and California. As the western Sharp-shinned Hawk occurs at Acapulco "only as a winter visitor," Mr. Nelson decides that for this reason the California bird "may be taken as typical of this form." Possibly a recognizable Northwest Coast form of the Sharp-shinned Hawk may yet be found, with a limited and fairly well defined breeding range, but until this has been made out neither of these names — *pacificus* from California and *rufilatus*, based on Fort Bridger specimens — need to give the layman any anxiety. In all probability neither will be available for the hypothetical new form, being apparently pure synonyms of *velox*, which seems to range across the continent without any satisfactorily recognizable western form, Rocky Mountain, Great Basin, and most California specimens, when comparable as to season and age, being not appreciably different from the eastern bird.

The common Turkey Buzzard of North America is shown to be separable from the Turkey Buzzard of Mexico, the West Indies and tropical America generally, through its much larger size and slight color difference. These were noticed by Wied in 1839, and for this reason he proposed the name *septentrionalis* for the North American bird (type locality, New Harmony, Indiana), thus restricting the name *aura* Linn. to the smaller southern form. The North American Turkey Buzzard, as Mr. Nelson shows, must stand as *Cathartes aura septentrionalis* (Wied). The same point is made, apparently independently and almost simultaneously, by Mr. Bangs in his 'Birds of the Isle of Pines' (Amer. Nat., April, 1905, p. 190, published April 26).

Mr. Nelson further shows that the Red-eyed Cowbird was first described by Lesson in 1839 as *Tangavius involucratus*, which name must replace the now current but much later *Collothrus robustus*. According to Mr. Nelson the three Mexican forms of *Tangavius* are merely subspecies and not species, so that the full name of our bird becomes *Tangavius æneus involucratus* (Lesson). — J. A. A.

Schiøler on the Greenland Mallard.¹ — On the basis of a comparison of a large series of specimens of the Mallard from Greenland with specimens from Denmark Mr. Schiøler has separated the Greenland form as a subspecies under the name *Anas boschas spilogaster*. The Greenland form differs from true *boschas* in being larger, with a somewhat shorter bill, darker upperparts and much more heavily spotted underparts. A large number of specimens are described in detail (including tables of measurements), and three plates, from photographs, very clearly illustrate the color differences claimed. — J. A. A.

¹Om dem grønlandske Stokand, *Anas boschas spilogaster*. Af E. Lehn Schiøler. Viedensk. Meddel. fra den naturh. Foren. i Kbhvn., 1905, pp. 129-148, pll. ii-iv.